arXiv:1602.06911v1 [math.AT] 22 Feb 2016

OBSTRUCTION THEORY FOR COINCIDENCES OF MULTIPLE MAPS

THAÍS MONIS AND PETER WONG

ABSTRACT. Let $f_1, ..., f_k : X \to N$ be maps from a complex X to a compact manifold N, $k \geq 2$. In previous works [1, 14], a Lefschetz type theorem was established so that the non-vanishing of a Lefschetz type coincidence class $L(f_1, ..., f_k)$ implies the existence of a coincidence $x \in X$ such that $f_1(x) = ... = f_k(x)$. In this paper, we investigate the converse of the Lefschetz coincidence theorem for multiple maps. In particular, we study the obstruction to deforming the maps $f_1, ..., f_k$ to be coincidence free. We construct an example of two maps $f_1, f_2 : M \to T$ from a sympletic 4-manifold M to the 2-torus T such that f_1 and f_2 cannot be homotopic to coincidence free maps but for *any* $f : M \to T$, the maps f_1, f_2, f are deformable to be coincidence free.

1. INTRODUCTION

The celebrated Lefschetz coincidence theorem states that for any two maps $f, g : M \to N$ between closed connected oriented triangulated n-manifolds, if the Lefschetz coincidence number (trace) $L(f, g)$ is non-zero then the coincidence set $C(f, g) = \{x \in M \mid f(x) = g(x)\}$ must be non-empty. However, the converse does not hold in general. In this direction, E. Fadell showed [4] that if N is simply-connected then the vanishing of $L(f, g)$ is sufficient to deform the maps $f \sim f', g \sim g'$ so that $C(f', g') = \emptyset$. For non-simply connected N, the vanishing of the Nielsen number $N(f, g)$ often provides the converse. Following [4] and [5], it was shown in [8] that the (primary) obstruction $o_n(f, g)$ $(n \geq 3)$ to deforming f and g to be coincidence free is Poincaré dual to the twisted Thom class of the normal bundle of the diagonal $\Delta(N)$ in $N \times N$ with appropriate local coefficients.

2010 *Mathematics Subject Classification.* Primary: 55M20; secondary: 55S35.

Key words and phrases. Obstruction theory, Lefschetz coincidence theory, local coefficients.

This work was initiated during the second author's visit to the Department of Mathematics at UNESP-Rio Claro, May 18 - June 17, 2014 and completed during his visit to the department Feb. 21 - 26, 2016. The second author would like to thank the Mathematics Department for the invitation and financial support. The first author was supported by FAPESP of Brazil Grant number 2014/17609-0.

Suppose M is a compact topological space, N is a closed connected oriented manifold, and $f_1, ..., f_k : M \to N$ are maps. A Lefschetz type coincidence class $L(f_1, ..., f_k)$ was defined in [1] and it was shown that $L(f_1, ..., f_k) \neq 0$ implies $C(f_1, ..., f_k) := \{x \in M \mid f_1(x) = ... = f_k(x)\}$ is non-empty. This Lefschetz type result has been extended to non-orientable N in [14].

The purpose of this paper is to examine the converse of this Lefschetz type theorem, that is, the problem when $f_1, ..., f_k$ can be deformed to be coincidence free. The approach here is via obstruction theory, following [4, 5, 2, 8]. We examine the primary obstruction to deforming $f_1, ..., f_k$ to be coincidence free on the $(k-1)n$-th skeleton where $n = \dim N$. We prove an analogous converse of the Lefschetz type coincidence theorem when N is simply-connected and $\dim M = (k-1)n$. We give further examples of Jiang-type spaces N for which the converse theorem holds. This paper is organized as follows. In section 2, we generalize the Lefschetz coincidence classes defined in [1] and in [14] to homomorphisms $\Lambda(f_1, ..., f_k; R_{N^k})$ with arbitrary local coefficients. This homomorphism is similar to a certain homomorphism ϕ of [9], relating to the preimage of a map. In section 3, we study the primary obstruction to deforming $f_1, ..., f_k$ to be coincidence free using the appropriate local coefficient system $\pi_j^*(F)$. The calculation of the local system $\pi_j^*(F)$ is carried out in section 4. We then compute the obstruction to deformation in section 5 and prove the converse of the Lefschetz coincidence theorem for multiple maps in section 6. We also compare our result to similar Nielsen type results of P. Staecker [16]. In section 7, we illustrate our results with examples. In particular, we construct two maps $f_1, f_2 : M \to N$, $\dim M > \dim N$ such that f_1, f_2 are not deformable to be coincidence free but for *any* $f : M \to N$, f_1, f_2, f are homotopic to be coincidence free (see Examples 7.1 and 7.2).

2. Lefschetz coincidence homomorphism with local coefficients

In this section, we review the Lefschetz coincidence classes introduced in [1] and in [14] and compare them with a more general homomorphism studied in [9].

Let X be a compact topological space and N be a connected closed n-manifold. For now, let us assume N is oriented. Suppose $f_1, ..., f_k : X \to N$ are maps, $\mu \in H^n(N \times N, N \times N \setminus \Delta(N))$ be the Thom class of the normal bundle of the diagonal $\Delta(N)$ in $N \times N$. Then, in [1], the authors defined

$$L_1(f_1, ..., f_k) = [(j \times ... \times j) \circ (h_1, ..., h_{k-1})]^*(\mu \times ... \times \mu) \tag{2.1}$$

where $h_i : X \to N \times N$ is given by $h_i(x) = (f_i(x), f_{i+1}(x))$ and $j : N \times N \hookrightarrow (N \times N, N \times N \setminus \Delta(N))$ is the inclusion. In [14], a similar Lefschetz type coincidence class was defined as follows. Let $\Delta_k(N) = \{(x, ..., x) \in N^k \mid x \in N\}$ be the diagonal of N in N^k and μ_k be the Thom class of the normal bundle of $\Delta_k(N)$ in N^k in $H^{(k-1)n}(N^k, N^k \setminus \Delta_k(N); R \times \Gamma_N^* \times ... \times \Gamma_N^*)$. Here R is a principal ideal domain and $\Gamma_N^* = \mathrm{Hom}(\Gamma_N, R)$ for the R-orientation system Γ_N on N. Define $\tilde{h}_i : X \to N$ by $\tilde{h}_i(x) = (f_1(x), f_{i+1}(x))$. Then

$$\begin{aligned} L_2(f_1, ..., f_k) &= [i \circ (f_1, ..., f_k)]^*(\mu_k) \\ &= [(j \times ... \times j) \circ (\tilde{h}_1, ..., \tilde{h}_{k-1})]^*(\mu \times ... \times \mu) \end{aligned} \tag{2.2}$$

where $i : N^k \to (N^k, N^k \setminus \Delta_k(N))$ is the inclusion. Note that

$$(j \times ... \times j) \circ (h_1, ..., h_{k-1}) = \sigma \circ i \circ (f_1, ..., f_k)$$

where $\sigma(x_1, ..., x_k) = ((x_1, x_2), (x_2, x_3), ..., (x_{k-1}, x_k))$. The induced homomorphism $\sigma^* : \otimes_{k-1}[H^n(N^2, N^2 \setminus \Delta(N))] \to H^{(k-1)n}(N^k, N^k \setminus \Delta_k(N)$ is an isomorphism when N is orientable. In this case, $\sigma^*(\mu \times ... \times \mu) = \mu_k$ and hence $L_1(f_1, ..., f_k) = L_2(f_1, ..., f_k)$.

A close inspection indicates that $L_2(f_1, ..., f_k)$ is in fact a special case of the coincidence index homomorphism studied in [9] which we recall as follows. Let X, Y be closed manifolds of dimension a and b, respectively. Suppose $B \subset Y$ is a closed submanifold of dimension ℓ. Let $F : X \to Y$ be transverse to B so that $C = F^{-1}(B)$ is a closed submanifold of dimension $a - b + \ell$. Suppose R_Y is a local system on Y with typical group R, $R_X = F^*(R_Y), R'_X = R_X \times \Gamma_X$ and $R'_C = i_1^*(R'_X)$ where $i_1 : C = F^{-1}(B) \hookrightarrow X$. For any $0 \leq j$, define $\phi(j)$ to be the composite homomorphism given by

$$H^j(Y, Y \setminus B; R_Y) \xrightarrow{F^*} H^j(X, X \setminus C; R_X) \xrightarrow{A^{-1}} H_{a+b-j}(C; R'_C)$$

where A represents the Alexander Duality isomorphism (see e.g. [15]). It was shown in [9, Theorem 2.4] that for any $r \in H^j(Y, Y \setminus B; R_Y)$,

$$i_{1*}\phi(j)(r) = F^*(j_2^*(r)) \cap [z_X]$$

where $j_2 : Y \hookrightarrow (Y, Y \setminus B)$ and $[z_X]$ is the (twisted) fundamental class of X with coefficients in R_X. It follows from sections 4.3 and 4.5 of [9] that $\phi((k-1)n) \neq 0 \Rightarrow C \neq \emptyset$. Now let $a = (k-1)n, b = kn, Y = N^k, B = \Delta_k(N), F = (f_1, ..., f_k), C = C(f_1, ..., f_k)$ and $j = (k-1)n$. If $R_Y = R \times \Gamma_N^* \times ... \times \Gamma_N^*$ then it is easy to see that

$$L_2(f_1, ..., f_k) = I^* \circ A \circ \phi((k-1)n)(\mu_k) \tag{2.3}$$

where $I : X \hookrightarrow (X, X \setminus C)$ is the inclusion. Now $L_2(f_1, ..., f_k) \neq 0$ implies that $\phi((k-1)n) \neq 0$. Thus, the Lefschetz type coincidence theorems of [14] and of [1] follow immediately.

Next, we define a Lefschetz coincidence homomorphism with any arbitrary local coefficient system R_{N^k}.

Let X be a compact topological space and N a closed manifold of dimension n. Suppose $f_1, ..., f_k : X \to N$ are maps. Let $F = (f_1, ..., f_k) : (X, X \setminus C(f_1, ..., f_k)) \to (N^k, N^k \setminus \Delta_k(N))$, $I : X \hookrightarrow (X, X - C(f_1, ..., f_k))$. Define the *Lefschetz coincidence homomorphism of* $f_1, ..., f_k$ *with local coefficients* R_{N^k} to be

(2.4)
$$\Lambda(f_1, ..., f_k; R_{N^k}) := I^* \circ F^* : H^{(k-1)n}(N^k, N^k \setminus \Delta_k(N); R_{N^k}) \to H^{(k-1)n}(X; F^*(R_{N^k})).$$

Thus, we have

$$\Lambda(f_1, ..., f_k; R \times \Gamma_N^* \times ... \times \Gamma_N^*)(\mu_k) := i^* \circ (f_1, ..., f_k)^*(\mu_k) = L_2(f_1, ..., f_k) \tag{2.5}$$

where $i : N^k \hookrightarrow (N^k, N^k \setminus \Delta_k(N))$. Now, the following result, which generalizes the main theorems of [1, 14], is immediate.

Theorem 2.1. *Let X be a compact topological space, N a closed manifold of dimension n and $k \geq 3$. For any maps $f_1, ..., f_k : X \to N$ and local coefficient R_{N^k}, if $\Lambda(f_1, ..., f_k; R_{N^k}) \neq 0$ then $C(f_1, ..., f_k) \neq \emptyset$.*

In section 5, we will generalize the Lefschetz coincidence classes of [1, 14] to $\Lambda(f_1, ..., f_k; R_{N^k})(\tilde{\mu}_k)$ for an approriate local coefficient and cohomology class $\tilde{\mu}_k$ which will be called the twisted Thom class of N^k.

3. Obstructions and local coefficients

In [4], E. Fadell showed that the classical Lefschetz coincidence class $L(f, g)$ coincides with the obstruction to deforming f and g to be coincidence free provided the target manifold is simply-connected. For non-simply connected manifolds, the obstruction class [5] necessarily involves certain local coefficient system. We now recall, following the treatment in [5] or in [2], the local coefficients employed in the obstruction to deformation.

Let $f : X \to Y$ be a map from a finite connected complex X to a closed connected manifold Y and $B \subset Y$ be a closed submanifold. First, we replace the inclusion $i : Y \setminus B \hookrightarrow Y$ by a fiber map $p : E \to Y$ where $E = \{(x, \omega) \mid x \in Y \setminus B, \omega \in Y^{[0,1]}, \omega(0) = x\}$ and $p(x, \omega) = \omega(1)$. Let the typical fiber be $F = p^{-1}(x_0) = \{(x, \omega) \mid \omega(1) = x_0\}$ for some $x_0 \in Y \setminus B$. It follows that $\pi_{j+1}(F; \overline{x_0}) \cong \pi_j(Y, Y \setminus B; x_0)$.

It is straightforward to verify that f is deformable into $Y \setminus B$, i.e., $f \sim f'$ so that $f'(X) \subseteq Y \setminus B$, if and only if f can be lifted to a map $\tilde{f} : X \to E$ such that $p \circ \tilde{f} = f$. Furthermore, f is deformable into $Y \setminus B$ if and only if the pullback fibration $q : f^*(p) \to X$ induced by f has a (global) section.

If $(Y, Y \setminus B)$ is $(m-1)$ connected. It follows from the classical obstruction theory for lifting (see e.g. [19]) that the primary obstruction to finding a section to q is given by a class $o_m(f) \in H^m(X; \pi^*_{m-1}(F))$ where $\pi^*_{m-1}(F)$ is the local system on X induced by $f : X \to Y$ from the local system $\pi_{m-1}(F)$ on Y. This class is the obstruction to deforming f to f' with $f'(X^{(m)}) \subset Y \setminus B$ where $X^{(m)}$ is the m-skeleton of X.

Now let $X = M, Y = N^k, B = \Delta_k(N)$ and $f = (f_1, ..., f_k)$ where $f_1, ..., f_k : M \to N$ are maps from a finite connected CW-complex M to a closed connected manifold N of dimension n, where $k \geq 2$. Next, we will study the obstruction to deforming the maps $f_1, ..., f_k$ to be coincidence free but first we must calculate the local coefficient system $\pi^*_{m-1}(F)$.

In this setting, we set

$$E = E(N^k, N^k \setminus \Delta_k(N)) = \{(y, \omega) \mid y \in N^k \setminus \Delta_k(N) \text{ and } \omega : I \to N^k \text{ is a path with } \omega(0) = y\}.$$

Let $p : E \to N^k$ be the fiber map given by $p(y, \omega) = \omega(1)$. If we take a point $y_0 = (y_0^1, \ldots, y_0^k) \in N^k$ the fiber over y_0, $F = p^{-1}(y_0)$, is given by

$$F = \{(\omega(0), \omega) \mid \omega(1) = y_0\}.$$

In words, the fiber is the space of paths in N^k which begins in $N^k \setminus \Delta_k(N)$ and end at y_0. Let $y_0 \in N^k \setminus \Delta_k(N)$ and denote by $\overline{y_0}$ the constant path $\overline{y_0}(t) = y_0$ for every t. There is an isomorphism

$$\pi_j(F; \overline{y_0}) \simeq \pi_{j+1}(N^k, N^k \setminus \Delta_k(N); y_0).$$

The above identification can be seen as the following: the element $[\alpha] \in \pi_j(F; \overline{y_0})$, where $\alpha : (S_j, e) \to (F, \overline{y_0})$, is mapped to $[\beta] \in \pi_{j+1}(N^k, N^k \setminus \Delta_k(N); y_0)$, where $\beta : (D^{j+1}, S^j, e) \to$

$(N^k, N^k \setminus \Delta_k(N), y_0)$ is given by

$$\beta(e + t(v - e)) = \alpha(z)(1 - t),$$

where $e = (1, 0, \ldots, 0)$ and $v \in S^j$.

Now, since $(N^k, N^k \setminus \Delta_k(N))$ is $(n(k-1)-1)$-connected, F is $(n(k-1)-2)$-connected. Therefore, if $3 \leq n(k-1)$ then $\pi_{(n(k-1)-1)}(F)$ forms a local system on N^k. Our next step is to describe such local system $\pi_{(n(k-1)-1)}(F)$ on N^k.

4. The local system $\pi_{(n(k-1)-1)}(F)$

As we saw in the last section, if we choose a base point $y_0 = (y_0^1, \ldots, y_0^k) \in N^k \setminus \Delta_k(N)$, we have a natural identification

$$\psi_{y_0} : \pi_{n(k-1)}(N^k, N^k \setminus \Delta_k(N); y_0) \simeq \pi_{n(k-1)-1}(F; \overline{y_0}),$$

where $F = p^{-1}(y_0)$ and $\overline{y_0}$ is the constant path at y_0. In fact, ψ is an isomorphism of local systems on $N^k \setminus \Delta_k(N)$. We now determine the action of

$$\pi_1(N^k \setminus \Delta_k(N); y_0) = \pi_1(N; y_0^1) \times \cdots \times \pi_1(N; y_0^k) = \underbrace{\pi \times \cdots \times \pi}_{k-times}$$

on

$$\pi_{n(k-1)}(N^k, N^k \setminus \Delta_k(N); y_0).$$

Let $\eta : \tilde{N} \to N$ be the universal cover of N. We assume that $y_0 = (y_0^1, \ldots, y_0^k)$ lies inside a small tubular neighborhood of $\Delta_k(N)$ and we consider the diagram

$$\begin{array}{ccc} \tilde{N} & \longleftarrow & (\tilde{N}^k, \tilde{N}^k \setminus \xi^{-1}(\Delta_k(N))) \\ \eta \downarrow & & \downarrow \xi \\ N & \longleftarrow & (N^k, N^k \setminus \Delta_k(N)) \end{array}$$

where $\xi = \underbrace{\eta \times \cdots \times \eta}_{k-times}$ and the horizontal maps are projections on the first coordinate. These horizontal maps are fibered pairs with fibers $(N^{k-1}, N^{k-1} \setminus \underbrace{(y_0^1, y_0^1, \ldots, y_0^1)}_{(k-1)-times})$ (lower horizontal

map) and $(\tilde{N}^{k-1}, \tilde{N}^{k-1} \setminus \underbrace{\eta^{-1}(y_0^1) \times \eta^{-1}(y_0^1) \times \cdots \times \eta^{-1}(y_0^1)}_{(k-1)-times})$ (upper horizontal map). Then, choosing $\tilde{y}_0 = (\tilde{y}_0^1, \ldots, \tilde{y}_0^k) \in \tilde{N}^k$, we have the isomorphisms

$$\begin{aligned}
&\pi_{n(k-1)}(\tilde{N}^{k-1}, \tilde{N}^{k-1} \setminus \eta^{-1}(y_0^1) \times \eta^{-1}(y_0^1) \times \cdots \times \eta^{-1}(y_0^1); (\tilde{y}_0^2, \ldots, \tilde{y}_0^k)) \\
&\simeq \pi_{n(k-1)}(\tilde{N}^k, \tilde{N}^k \setminus \xi^{-1}(\Delta_k(N)); \tilde{y}_0)
\end{aligned}$$

and

$$\begin{aligned}
&\pi_{n(k-1)}(N^{k-1}, N^{k-1} \setminus (y_0^1, y_0^1, \ldots, y_0^1); (y_0^2, \ldots, y_0^k)) \\
&\simeq \pi_{n(k-1)}(N^k, N^k \setminus \Delta_k(N); y_0).
\end{aligned}$$

Moreover, because $\xi : (\tilde{N}^k, \tilde{N}^k \setminus \xi^{-1}(\Delta_k(N))) \to (N^k, N^k \setminus \Delta_k(N))$ is a covering map, we have

$$\pi_{n(k-1)}(\tilde{N}^k, \tilde{N}^k \setminus \xi^{-1}(\Delta_k(N)); \tilde{y}_0) \simeq \pi_{n(k-1)}(N^k, N^k \setminus \Delta_k(N); y_0).$$

Therefore

$$\begin{aligned}
&\pi_{n(k-1)}(N^k, N^k \setminus \Delta_k(N); y_0) \\
&\simeq \pi_{n(k-1)}(\tilde{N}^{k-1}, \tilde{N}^{k-1} \setminus \eta^{-1}(y_0^1) \times \eta^{-1}(y_0^1) \times \cdots \times \eta^{-1}(y_0^1); (\tilde{y}_0^2, \ldots, \tilde{y}_0^k)).
\end{aligned}$$

Furthermore, by simple connectivity, it follows from the relative Hurewicz theorem that

$$\begin{aligned}
&\pi_{n(k-1)}(\tilde{N}^{k-1}, \tilde{N}^{k-1} \setminus \eta^{-1}(y_0^1) \times \eta^{-1}(y_0^1) \times \cdots \times \eta^{-1}(y_0^1); (\tilde{y}_0^2, \ldots, \tilde{y}_0^k)) \\
&\simeq H_{n(k-1)}(\tilde{N}^{k-1}, \tilde{N}^{k-1} \setminus \eta^{-1}(y_0^1) \times \eta^{-1}(y_0^1) \times \cdots \times \eta^{-1}(y_0^1)).
\end{aligned}$$

Note that

$$(\tilde{N}^{k-1}, \tilde{N}^{k-1} \setminus \eta^{-1}(y_0^1) \times \eta^{-1}(y_0^1) \times \cdots \times \eta^{-1}(y_0^1)) = \underbrace{(\tilde{N}, \tilde{N} \setminus \eta^{-1}(y_0^1)) \times \cdots \times (\tilde{N}, \tilde{N} \setminus \eta^{-1}(y_0^1))}_{(k-1)-times}$$

and, therefore,

$$\begin{aligned}
&\pi_{n(k-1)}(N^k, N^k \setminus \Delta_k(N); y_0) \\
&\simeq \underbrace{H_n(\tilde{N}, \tilde{N} \setminus \eta^{-1}(y_0^1)) \otimes \cdots \otimes H_n(\tilde{N}, \tilde{N} \setminus \eta^{-1}(y_0^1))}_{(k-1)-times}.
\end{aligned}$$

Let $\eta^{-1}(y_0^1) = \{\tilde{y}_i\}_i$ with the convention that $\tilde{y}_1 = \tilde{y}_0^1$. Let V be a euclidean neighborhood of y_0^1 in N and $\tilde{V}_i$ euclidean neighborhood of $\tilde{y}_i$ such that $\eta^{-1}(V) = \sqcup_i \tilde{V}_i$ and $\eta : \tilde{V}_i \to V$ a

homeomorphism. By excision,

$$H_n(\tilde{N}, \tilde{N} \setminus \eta^{-1}(y_0^1)) \simeq \sum_i H_n(\tilde{V}_i, \tilde{V}_i \setminus \tilde{y}_i).$$

Identifying π with the covering transformations of $\tilde{N}$, given $\sigma \in \pi$, if $\sigma \tilde{y}_0^1 = \tilde{y}_i$ then $\sigma \tilde{V}_1 = \tilde{V}_i$.

Choose a local orientation of V at y_0^1, which determines a generator $\gamma_1 \in H_n(\tilde{V}_1, \tilde{V}_1 \setminus \tilde{y}_0^1)$. Let $\alpha \in \pi$ and set $\gamma_\alpha = \alpha\gamma_1$. Thus, γ_α generates $H_n(\tilde{V}_i, \tilde{V}_i \setminus \tilde{y}_i)$ if $\alpha \tilde{y}_0^1 = \tilde{y}_i$.

Since $(\tilde{N}^{k-1}, \tilde{N}^{k-1} \setminus \eta^{-1}(y_0^1) \times \eta^{-1}(y_0^1) \times \cdots \times \eta^{-1}(y_0^1))$ is the fiber of the fiber pair map

$$\tilde{N} \leftarrow (\tilde{N}^k, \tilde{N}^k \setminus \xi^{-1}(\Delta_k(N)))$$

over each $\tilde{y}_i$, for each $\tilde{y}_i$ we have a fiber inclusion

$$\theta_i : (\tilde{N}^{k-1}, \tilde{N}^{k-1} \setminus \eta^{-1}(y_0^1) \times \eta^{-1}(y_0^1) \times \cdots \times \eta^{-1}(y_0^1)) \hookrightarrow (\tilde{N}^k, \tilde{N}^k \setminus \xi^{-1}(\Delta_k(N)))$$

given by

$$\theta_i(u) = (\tilde{y}_i, u), \ u \in \tilde{N}^{k-1}.$$

Now, we identify $\mathbb{Z}[\underbrace{\pi \times \cdots \times \pi}_{(k-1)-times}]$ with the image of

$$\underbrace{H_n(\tilde{N}, \tilde{N} \setminus \eta^{-1}(y_0^1)) \otimes \cdots \otimes H_n(\tilde{N}, \tilde{N} \setminus \eta^{-1}(y_0^1))}_{(k-1)-times}$$

under θ_{1*} via the correspondence

$$(\alpha_1, \ldots, \alpha_{k-1}) \mapsto \theta_{1*}(\gamma_{\alpha_1} \otimes \cdots \otimes \gamma_{\alpha_{k-1}}) = \theta_{1*}(\alpha_1\gamma_1 \otimes \cdots \otimes \alpha_{k-1}\gamma_1).$$

Observe that $\theta_{1*}(\gamma_1 \otimes \cdots \otimes \gamma_1)$ can be represented by an $n(k-1)$-cell $\{\tilde{y}_0^1\} \times (D_1^n)^{k-1}$ in $\tilde{N}^k$ transverse to $\xi^{-1}(\Delta_k(N))$ at $(\tilde{y}_0^1, \tilde{y}_0^1, \ldots, \tilde{y}_0^1)$. Thus, for $\sigma \in \pi$, the diagonal element

$$(\sigma, \sigma, \ldots, \sigma) \in \underbrace{\pi \times \pi \times \cdots \times \pi}_{k-times}$$

sends $\{\tilde{y}_0^1\} \times (D_1^n)^{k-1}$ to an $n(k-1)$-cell $\{\tilde{y}_i\} \times (D_i^n)^{k-1}$ transverse to $\xi^{-1}(\Delta_k(N))$ at $(\tilde{y}_i, \tilde{y}_i, \ldots, \tilde{y}_i)$ if $\sigma \tilde{y}_0^1 = \tilde{y}_i$. Thus, one can see that

$$(\sigma, \sigma, \ldots, \sigma)\theta_{1*}(\gamma_1 \otimes \gamma_1 \otimes \cdots \otimes \gamma_1) = (\operatorname{sgn}\sigma)^{k-1}\theta_{1*}(\gamma_1 \otimes \gamma_1 \otimes \cdots \otimes \gamma_1).$$

Now, we can compute the action of π^k on $\mathbb{Z}[\pi^{k-1}]$ via the above identifications:

(4.1)
$$\begin{aligned}(\sigma_1,\dots,\sigma_k)\circ(\alpha_1,\dots,\alpha_{k-1}) &\equiv (\sigma_1,\dots,\sigma_k)\theta_{1*}(\alpha_1\gamma_1\otimes\cdots\otimes\alpha_{k-1}\gamma_1)\\ &= (\sigma_1,\dots,\sigma_k)(1\times\alpha_1\times\cdots\times\alpha_{k-1})\theta_{1*}(\gamma_1\otimes\cdots\otimes\gamma_1)\\ &= (\sigma_1,\sigma_2\alpha_1,\dots,\sigma_k\alpha_{k-1})\theta_{1*}(\gamma_1\otimes\cdots\otimes\gamma_1)\\ &= (1,\sigma_2\alpha_1\sigma_1^{-1},\dots,\sigma_k\alpha_{k-1}\sigma_1^{-1})(\sigma_1,\dots,\sigma_1)\theta_{1*}(\gamma_1\otimes\cdots\otimes\gamma_1)\\ &= (\operatorname{sgn}\sigma_1)^{k-1}(1,\sigma_2\alpha_1\sigma_1^{-1},\dots,\sigma_k\alpha_{k-1}\sigma_1^{-1})\theta_{1*}(\gamma_1\otimes\cdots\otimes\gamma_1)\\ &= (\operatorname{sgn}\sigma_1)^{k-1}\theta_{1*}(\sigma_2\alpha_1\sigma_1^{-1}\gamma_1\otimes\cdots\otimes\sigma_k\alpha_{k-1}\sigma_1^{-1}\gamma_1)\\ &\equiv (\operatorname{sgn}\sigma_1)^{k-1}(\sigma_2\alpha_1\sigma_1^{-1},\dots,\sigma_k\alpha_{k-1}\sigma_1^{-1}).\end{aligned}$$

By pulling back the local system by the map $f: M\to N^k$, the action of $\pi_1(M)$ on $\mathbb{Z}[\pi^{k-1}]$ is given by

(4.2)
$$\begin{aligned}\gamma\circ(\alpha_1,\dots,\alpha_{k-1}) &= (\varphi_1(\gamma),...,\varphi_k(\gamma))\circ(\alpha_1,\dots,\alpha_{k-1})\\ &= (\operatorname{sgn}\varphi_1(\gamma))^{k-1}(\varphi_2(\gamma)\alpha_1\varphi_1(\gamma)^{-1},\dots,\varphi_k(\gamma)\alpha_{k-1}\varphi_1(\gamma)^{-1}).\end{aligned}$$

Here, φ_i is the homomorphism induced by f_i.

5. Lefschetz coincidence class as primary obstruction

Following [5], the primary obstruction to deforming f off a subspace B was defined as a *universal element* in [2]. In our setting, first we triangulate N and hence N^k so that the $kn-1$ skeleton $(N^k)^{(kn-1)}\subset N^k\setminus\Delta_k(N)$ for $k\geq 3$. Then there is the (only) primary obstruction $\tilde{\mu}_k\in H^{(k-1)n}(N^k,N^k\setminus\Delta_k(N);\mathbb{Z}[\pi^{k-1}])$ to deforming the identity map 1_{N^k} off the subspace $\Delta_k(N)$. We call this element $\tilde{\mu}_k$ the *twisted Thom class* of the normal bundle of $\Delta_k(N)$ in N^k or simply the *twisted Thom class* of N^k.

For any i,j, consider the map

$$e:(N^{i+j-1},N^{i+j-1}\setminus\Delta_{i+j-1}(N))\to(N^i,N^i\setminus\Delta_i(N))\times(N^j,N^j\setminus\Delta_j(N))$$

given by $e(x_1,x_2,...,x_{i+j-1})=((x_1,x_2,...,x_i),(x_1,x_{i+1},x_{i+2},...,x_{i+j-1})$.

Proposition 5.1. The map e induces a homomorphism

$$H^{(i-1)n}(N^{\times_i};\mathbb{Z}[\pi^{i-1}])\otimes H^{(j-1)n}(N^{\times_j};\mathbb{Z}[\pi^{j-1}])\to H^{(i+j-2)n}(N^{\times_{i+j-1}};\mathbb{Z}[\pi^{i+j-2}])$$

such that $e^*(\tilde{\mu}_i\otimes\tilde{\mu}_j)=\tilde{\mu}_{i+j-1}$. Here $N^{\times_k}$ denotes the pair $(N^k,N^k\setminus\Delta_k(N))$ and $\pi=\pi_1(N)$.

Proof. First there is an Eilenberg-Zilber map

$$\mathrm{EZ} : C_*(N^{\times_i}; \mathbb{Z}[\pi^{i-1}]) \otimes C_*(N^{\times_j}; \mathbb{Z}[\pi^{j-1}]) \to C_*(N^{\times_{i+j}}; \mathbb{Z}[\pi^{i-1}] \otimes \mathbb{Z}[\pi^{j-1}])$$

where the action of $\pi^{i+j} = \pi^i \times \pi^j$ on $\mathbb{Z}[\pi^{i-1}] \otimes \mathbb{Z}[\pi^{j-1}]$ is the diagonal action. More precisely, for $(\sigma_1, ..., \sigma_{i+j}) \in \pi^{i+j}$, we have

$$\begin{aligned} &(\sigma_1, ..., \sigma_{i+j}) \circ (\alpha_1, ..., \alpha_{i-1}, \beta_1, ..., \beta_{j-1}) \\ =&(\mathrm{sgn}\sigma_1)^{i-1}(\sigma_2\alpha_1\sigma_1^{-1}, ..., \sigma_i\alpha_{i-1}\sigma_1^{-1}) \otimes (\mathrm{sgn}\sigma_{i+1})^{j-1}(\sigma_{i+2}\beta_1\sigma_{i+1}^{-1}, ..., \sigma_{i+j}\beta_{j-1}\sigma_{i+1}^{-1}). \end{aligned} \tag{5.1}$$

Now the map e induces a local system $e^*(\mathbb{Z}[\pi^{i-1}] \otimes \mathbb{Z}[\pi^{j-1}])$ on $N^{\times_{i+j-1}}$ and this system is given by the following action:

$$\begin{aligned} &(\sigma_1, ..., \sigma_{i+j-1}) * (\alpha_1, ..., \alpha_{i-1}, \beta_1, ..., \beta_{j-1}) \\ =&e_\#(\sigma_1, ..., \sigma_{i+j-1}) \circ (\alpha_1, ..., \alpha_{i-1}, \beta_1, ..., \beta_{j-1}) \\ =&(\sigma_1, ..., \sigma_i, \sigma_1, \sigma_{i+1}, ..., \sigma_{i+j-1}) \circ (\alpha_1, ..., \alpha_{i-1}, \beta_1, ..., \beta_{j-1}) \\ =&(\mathrm{sgn}\sigma_1)^{i-1}(\sigma_2\alpha_1\sigma_1^{-1}, ..., \sigma_i\alpha_{i-1}\sigma_1^{-1}) \cdot (\mathrm{sgn}\sigma_1)^{j-1}(\sigma_{i+1}\beta_1\sigma_1^{-1}, ..., \sigma_{i+j-1}\beta_{j-1}\sigma_1^{-1}) \end{aligned}$$

This action coincides with that of (4.1) so that $e^*(\mathbb{Z}[\pi^{i-1}] \otimes \mathbb{Z}[\pi^{j-1}])$ coincides with the local system $\mathbb{Z}[\pi^{i+j-2}]$ discussed above. Now the obstruction to deforming the identity map 1_{N^q} off the subspace $\Delta_q(N)$ has a simple cochain representation given by $c_{(q-1)n}(1_{N^q})(\sigma) = [\sigma] \in \pi_{(q-1)n}(N^{\times_q})$. Let $\sigma_1 \times \sigma_2 \times ... \times \sigma_{i+j-1}$ be an $(i+j-1)n$ simplex where each σ_i is an n-simplex in N and $\tilde{c}_p$ denote the cochain representing the twisted Thom class $\tilde{\mu}_p$. Then

$$\begin{aligned} \langle \tilde{c}_{i+j-1}, \sigma_1 \times \sigma_2 \times ... \times \sigma_{i+j-1} \rangle &= [\sigma_1 \times \sigma_2 \times ... \times \sigma_i \times \sigma_{i+1} \times ... \times \sigma_{i+j-1}] \qquad \in \mathbb{Z}[\pi^{i+j-2}] \\ &= e^*([\sigma_1 \times \sigma_2 \times ... \times \sigma_i] \otimes [\sigma_1 \times \sigma_{i+1} \times ... \times \sigma_{i+j-1}]) \\ &= e^*(\langle \tilde{c}_i, \sigma_1 \times \sigma_2 \times ... \times \sigma_i \rangle \otimes \langle \tilde{c}_j, \sigma_1 \times \sigma_{i+1} \times ... \times \sigma_{i+j-1}] \rangle) \\ &= e^*(\langle \tilde{c}_i \otimes \tilde{c}_j, (\sigma_1 \times \sigma_2 \times ... \times \sigma_i) \otimes (\sigma_1 \times \sigma_{i+1} \times ... \times \sigma_{i+j-1}) \rangle) \\ &= \langle \tilde{c}_i \otimes \tilde{c}_j, e_*((\sigma_1 \times \sigma_2 \times ... \times \sigma_i) \otimes (\sigma_1 \times \sigma_{i+1} \times ... \times \sigma_{i+j-1})) \rangle. \end{aligned}$$

It follows that $e^*(\tilde{\mu}_i \otimes \tilde{\mu}_j) = \tilde{\mu}_{i+j-1}$. □

As an immediate corollary, we have the following useful result.

Corollary 5.2. Let $e' : N^{\times_k} \to \underbrace{N^{\times_2} \times ... \times N^{\times_2}}_{k-1}$ be defined by

$$e'(x_1, ..., x_k) = ((x_1, x_2), (x_1, x_3), ..., (x_1, x_k)).$$

Suppose $n = \dim N \geq 3$. Then e' induces a homomorphism e'^* such that $e'^*(\underbrace{\tilde{\mu}_2 \otimes ... \otimes \tilde{\mu}_2}_{k-1}) = \tilde{\mu}_k$.

Using the twisted Thom class as an element in the cohomology of $N^{\times_k}$ with local coefficients $\mathbb{Z}[\pi^{k-1}]$, we define the *twisted Lefschetz coincidence class of* $f_1, ..., f_k$ to be the element $\mathcal{L}(f_1, ..., f_k) := \Lambda(f_1, ..., f_k; \mathbb{Z}[\pi^{k-1}])(\tilde{\mu}_k)$. Since $\tilde{\mu}_k$ is the obstruction to deforming the identity off the subspace $\Delta_k(N)$, it follows from [2] that $\mathcal{L}(f_1, ..., f_k) = o_{(k-1)n}(f_1, ..., f_k)$ the primary obstruction to deforming $f_1, ..., f_k$ to be coincidence free on the $(k-1)n$ skeleton of X.

In the case when N is oriented, it was already shown in [8] that the primary obstruction to deforming f and g is mapped to the classical Lefschetz coincidence number $L(f, g)$ under the augmentation homomorphism $\mathbb{Z}[\pi] \to \mathbb{Z}$. Next, we show an analogous result, that is, there is a natural homomorphism $\tilde{\eta}$ induced by the augmentation map such that $\tilde{\eta}(\mathcal{L}(f_1, ..., f_k)) = L_2(f_1, ..., f_k)$. Here, we assume the principal ideal domain R is $\mathbb{Z}$.

Theorem 5.3. *The augmentation map* $\epsilon^{k-1} : \mathbb{Z}[\pi^{k-1}] \to \Gamma_N^* \times ... \times \Gamma_N^*$ *induces a homomorphism* $\tilde{\eta}$ *such that* $\tilde{\eta}(\mathcal{L}(f_1, ..., f_k)) = L_2(f_1, ..., f_k)$.

Proof. By the Alexander duality isomorphism (see [15, Thm. 6.4]), we have the following commutative diagram.

$$\begin{array}{ccc} H^{(k-1)n}(N^{\times_k}; \mathbb{Z}[\pi^{k-1}]) & \xrightarrow{A_1^{-1}} & H_n(\Delta_k(N); \Gamma_{N^k} \otimes \mathbb{Z}[\pi^{k-1}]) \\ \downarrow{\scriptstyle \eta} & & \downarrow{\scriptstyle 1 \otimes \epsilon^{k-1}} \\ H^{(k-1)n}(N^{\times_k}; \mathbb{Z} \times \Gamma_N^* \times ... \times \Gamma_N^*) & \xleftarrow{A_2} & H_n(\Delta_k(N); \Gamma_{N^k} \otimes \mathbb{Z} \times \Gamma_N^* \times ... \times \Gamma_N^*) \end{array}$$

Here, A_1 and A_2 are the corresponding duality isomorphisms so that $\eta = A_2 \circ (1 \otimes \epsilon^{k-1}) \circ A_1^{-1}$ where ϵ^{k-1} is the augmentation map. For any $\sigma \in \pi$, the action of $(\sigma, ..., \sigma) \in \pi_1(\Delta_k(N))$ on $\mathbb{Z} \times \Gamma_N^* \times ... \times \Gamma_N^*$ is the same as the action on $\mathbb{Z}[\pi^{k-1}]$ followed by ϵ^{k-1}. This implies that the twisted fundamental class of $\Delta_k(N)$ with coefficients in $\Gamma_{N^k} \otimes \mathbb{Z}[\pi^{k-1}]$ is mapped under $1 \otimes \epsilon^{k-1}$ to the twisted fundamental class of $\Delta_k(N)$ with coefficients in $\Gamma_{N^k} \otimes \mathbb{Z} \times \Gamma_N^* \times ... \times \Gamma_N^*$. By duality, the twisted Thom class $\tilde{\mu}_k$ is mapped under η to μ_k. Define $\tilde{\eta}$ by $\langle \tilde{\eta}(m), \sigma \rangle := \eta(\langle m, \sigma \rangle)$. Pulling back the classes $\tilde{\mu}_k, \mu_k$ yields the assertion. □

Theorem 5.4. *Let* $f_1, ..., f_k : X \to N$ *be maps from a finite complex* X *to a closed connected manifold* N *of dimension* n, $n \geq 3$. *Then*

$$o_{(k-1)n}(f_1, ..., f_k) = o_n(f_1, f_2) \cup o_n(f_1, f_3) \cup ... \cup o_n(f_1, f_k).$$

Proof. In Corollary 5.2 we state that if $e' : N^{\times_k} \to \underbrace{N^{\times_2} \times ... \times N^{\times_2}}_{k-1}$ is the map defined by $e'(x_1, ..., x_k) = ((x_1, x_2), (x_1, x_3), ..., (x_1, x_k))$ then $e'^*(\underbrace{\tilde{\mu}_2 \otimes ... \otimes \tilde{\mu}_2}_{k-1}) = \tilde{\mu}_k$. Let us consider the inclusions $I : X \to (X, X \setminus C(f_1, \ldots, f_k))$ and $j_i : X \to (X, X \setminus C(f_1, f_i))$, $i = 2, \ldots, k$. Then,

$$o_{(k-1)n}(f_1, ..., f_k) = I^*(f_1, \ldots, f_k)^*(\tilde{\mu}_k)$$

and

$$o_n(f_1, f_i) = j_i^*(f_1, f_i)^*(\tilde{\mu}_2),\ i = 2, \ldots, k.$$

Note that $e' \circ (f_1, \ldots, f_k) \circ I = ((f_1, f_2) \times \ldots \times (f_1, f_k)) \circ (j_1, \ldots, j_{k-1})$. Therefore

$$\begin{aligned} o_{(k-1)n}(f_1, ..., f_k) &= I^*(f_1, \ldots, f_k)^*(\tilde{\mu}_k) \\ &= I^*(f_1, \ldots, f_k)^*(e'^*(\tilde{\mu}_2 \otimes ... \otimes \tilde{\mu}_2)) \\ &= (j_1, \ldots, j_{k-1})^* \circ ((f_1, f_2) \times \ldots \times (f_1, f_k))^*(\tilde{\mu}_2 \otimes ... \otimes \tilde{\mu}_2) \\ &= j_1^*(f_1, f_2)^*(\tilde{\mu}_2) \cup j_2^*(f_1, f_3)^*(\tilde{\mu}_2) \cup ... \cup j_{k-1}^*(f_1, f_k)^*(\tilde{\mu}_2) \\ &= o_n(f_1, f_2) \cup o_n(f_1, f_3) \cup ... \cup o_n(f_1, f_k) \end{aligned}$$

□

Remark 5.5. Since $o_{(k-1)n}(f_1, ..., f_k) = \mathcal{L}(f_1, ..., f_k)$ and $o_n(f_1, f_i) = \mathcal{L}(f_1, f_i)$, Theorem 5.4 states that for $n \geq 3$, we have

$$\mathcal{L}(f_1, ..., f_k) = \mathcal{L}(f_1, f_2) \cup \mathcal{L}(f_1, f_3) \cup ... \cup \mathcal{L}(f_1, f_k). \tag{5.2}$$

Combining with the homomorphism $\tilde{\eta}$ as in Theorem 5.3, it is easy to establish the following product

$$L_2(f_1, ..., f_k) = L_2(f_1, f_2) \cup L_2(f_1, f_3) \cup ... \cup L_2(f_1, f_k)$$

which is obtained in [14, Thm. 4.2].

6. Converse of Lefschetz Coincidence Theorem

The converse of the Lefschetz coincidence theorem in our setting amounts to studying the problem when the maps $f_1, ..., f_k$ can be deformed to be coincidence free. For instance in the classical case, it has been shown [4] that when N is simply-connected then the vanishing of the Lefschetz coincidence number provides a converse to the Lefschetz coincidence theorem.

To study this problem for multiple maps, we use the obstruction theory developed in the previous sections.

Next, we need to calculate the obstruction cocycle, following [2, pp.20-21]. Suppose M is a closed oriented $(k-1)n$-dimensional manifold. By general position, we may assume that the map $f = (f_1, ..., f_k) : M \to N^k$ is transverse to the submanifold $\Delta_k(N)$ so that $C = f^{-1}(\Delta_k(N)) = C(f_1, ..., f_k)$ is a finite set of coincidence points of $f_1, \ldots, f_k$. Let $C = \{s_1, ..., s_r\}$. For each isolated coincidence point s_i, we may assume that there is a maximal $(k-1)n$-simplex σ_i containing s_i such that $C \cap \overline{\sigma}_i = \{s_i\}$. Following [5, 2], the primary obstruction cocycle $c_{(k-1)n}(f)$ to deforming f into the subspace $N^k \setminus \Delta_k(N)$ is given by

$$c_{(k-1)n}(f)(\sigma) = \begin{cases} 0 & \text{if } \sigma \cap C = \emptyset \\ \lambda(f, s_i)[f|_\sigma] & \text{if } \sigma = \sigma_i. \end{cases} \tag{6.1}$$

Here, $f|_{\sigma_i}$ represents the restriction map from $(\sigma_i, \partial\sigma_i)$ to $(V_i \times ... \times V_i, V_i \times ... \times V_i \setminus \{(y_i, ..., y_i)\})$ where V_i is an open neighborhood of $y_i = f_1(s_i) = ... = f_k(s_i)$ in N. Thus, $[f|_{\sigma_i}] \in \pi_{(k-1)n}(N^k, N^k \setminus \Delta_k(N)) \cong \pi_{(k-1)n-1}(F)$. Furthermore the coefficient $\lambda(f, s_i) = f|^*_{\sigma_i}(\tilde{\mu}_k)$ is the *local coincidence index* defined by the restriction $f|_{\sigma_i}$ using the homomorphism

$$f|^*_{\sigma_i} : H^{(k-1)n}(N^k, N^k \setminus \Delta_k(N)) \to H^{(k-1)n}(\sigma_i, \sigma_i \setminus \{s_i\}).$$

Since $(M, M - C) = (M, M - \{s_1, ..., s_r\})$, it follows from excision and additivity that

$$I^* \circ f^*(\tilde{\mu}_k) = \sum_{i=1}^{r} \lambda(f, s_i)$$

where $I : M \hookrightarrow (M, M - C)$. It follows that

$$c_{(k-1)n}(f) = \sum_{i}^{r} \lambda(f, s_i)[f|_{\sigma_i}] \tag{6.2}$$

and

$$\mathcal{L}(f_1, ..., f_k) = \sum_{i=1}^{r} \lambda(f, s_i). \tag{6.3}$$

Theorem 6.1. *Let $f_1, \ldots, f_k : M \to N$ from a closed connected $(k-1)n$-manifold M to a closed connected n-manifold N. Then $f_1, \ldots, f_k$ are deformable to be coincidence free if, and only if, $o_{(k-1)n}(f_1, ..., f_k) = 0$ where the obstruction cocycle is given by* (6.2).

In the case where N is simply-connected, $o_{(k-1)n}(f_1,...,f_k) = \mathcal{L}(f_1,...,f_k) = L_2(f_1,...,f_k) = L_1(f_1,...,f_k)$ since N is orientable and the coefficients are simple. In particular, when $N = S^n$ is the n-sphere, the formula of [1, Prop. 3.2] becomes

$$o_{(k-1)n}(f_1,...,f_n) = \sum_{i=0}^{k-1}(-1)^{(i-1)n}\widehat{f_{k-i}}^*(e)$$

where $e = \underbrace{e_n \times ... \times e_n}_{(k-1)-times}$, e_n the cohomology fundamental class of S^n, $\widehat{f_j} = (f_1,...,f_{j-1},f_{j+1},...,f_k)$. This formula generalizes a similar formula in [8, Theorem 3.4]. Furthermore, the equality $o_{(k-1)n}(f_1,...,f_k) = L_2(f_1,\ldots,f_k)$ generalizes [4, Theorem 1.1]) and we have the following converse theorem of the Lefschetz coincidence theorem when N is simply connected.

Theorem 6.2. *Let $f_1,\ldots,f_k : M \to N$ from a closed connected $(k-1)n$-manifold M to a closed connected and simply-connected n-manifold N. Then $f_1,\ldots,f_k$ are deformable to be coincidence free if, and only if, $L_2(f_1,...,f_k) = 0$.*

To obtain the converse Lefschetz theorem, Nielsen coincidence theory is often employed. In [16], P. Staecker independently investigated the coincidence problem for multiple maps from the Nielsen coincidence theory point of view. More precisely, Staecker considered the following two maps $F = (f_1,...,f_1), G = (f_2,...,f_k) : M \to N^{k-1}$ and employed the usual Nielsen coincidence theory by defining the Lefschetz and Nielsen coincidence numbers $L(F,G)$ and $N(F,G)$. When N is non-orientable, an appropriate (semi)index was used to define essentiality of the coincidence classes. We should point out that when N is orientable, $|L_2(f_1,...,f_k)| = |L(F,G)|$ and $o_{(k-1)n}(f_1,...,f_k) = 0 \Leftrightarrow N(F,G) = 0$. In this case, since $C = C(f_1,...,f_k) = \{s_1,...,s_r\}$ is a finite set, C is a disjoint union of $N(F,G)$ coincidence classes. Thus, the coincidence classes will have index of the same sign, if N is a Jiang space; a nilmanifold [11], an orientable coset space G/K of a compact connected Lie group G by a closed subgroup K [18]; or a $\mathcal{C}$-nilpotent space whose fundamental group has a finite index center [12] where $\mathcal{C}$ denotes the class of finite groups. For these spaces, $L(F,G) = 0 \Rightarrow N(F,G) = 0$. Thus, we have the following converse theorem.

Theorem 6.3. *Let $f_1,\ldots,f_k : M \to N$ from a closed connected $(k-1)n$-manifold M to a closed connected orientable n-manifold N. Suppose N is a Jiang space; a nilmanifold, an orientable coset space G/K of a compact connected Lie group G by a closed subgroup K; or a $\mathcal{C}$-nilpotent space whose fundamental group has a finite index center where $\mathcal{C}$ is the*

class of finite groups. Then $f_1, \ldots, f_k$ are deformable to be coincidence free if, and only if, $L_2(f_1, ..., f_k) = 0$.

7. Examples

When $n = 2$ and $k \geq 3$, $(k-1)n \geq 3$ so that $\pi_{(k-1)n-1}(F) \cong \pi_{(k-1)n}(N^k, N^k \setminus \Delta_k(N))$ is an abelian group so it is a local coefficient system on N^k even when N is a surface. Thus, Theorem 6.2 and Theorem 6.3 are valid when $n = 2$ provided $k \geq 3$. Next we illustrate by an example in which $f_1, f_2, f_3 : X \to S_g$ are maps from a closed 4-manifold to a surface S_g such that f_1, f_i cannot be deformed to be coincidence free for $i = 2, 3$. However, f_1, f_2, f_3 are deformable to be coincidence free.

Example 7.1. Let $X = S^1 \times S^3$, $f_1 : X \to S^2$ be given by $f_1(x_1, x_2) = h(x_2)$ where $x_2 \in S^3$ and $h : S^3 \to S^2$ is the Hopf map. Let $f_2 = \overline{c}$ be the constant map at some $c \in S^2$. Note that $f_1 : X \to S^2$ is a fibration and $f_1^*([\mu]) = 0$ where $[\mu]$ is the cohomology fundamental group of the base space S^2. This follows from the fact that $H^2(X) = 0$. Thus, we conclude that $L_2(f_1, f_2) = 0$. Note that f_1 and f_2 cannot be deformed to be coincidence free. To see that, let us assume that $f_1 \sim f_1'$ such that $f_1'(X) \subset S^2 - \{c\}$. It follows that $f_1'(X)$ is a subset of the complement of some open disk around c. This complement is homeomorphic to a closed 2-disk which is contractible, we conclude that f_1' is null-homotopic which in turn implies that the Hopf map would be null-homotopic, a contracdiction. Now let $f_3 : X \to S^2$ be *any* map. Since $L_2(f_1, f_2) = 0$, by [14, Thm. 4.2], we have $L_2(f_1, f_2, f_3) = 0$. The 2-sphere S^2 is simply connected so by Theorem 6.2, f_1, f_2, f_3 are deformable to be coincidence free. In particular, if we choose $f_3 = f_2$ then f_1, f_i are not deformable to be coincidence free for $i = 2, 3$. Note that X is not symplectic so by the classical condition of Thurston [17], the fiber $S^1 \times S^1$ is null homologous in X. As it turns out, Thurston's condition is only necessary for X to be symplectic as we use that to contruct the next example.

Example 7.2. According to R. Geiges [6], there exist symplectic 4-manifolds X that is a principal T^2-bundle over T^2 but the fibration is not symplectic so that $i_*([F]) = 0$ in $H_2(X)$ where $[F]$ is the fundamental class of the fiber $F = T^2$. According to a result of Gottlieb (see its generalization in [9]), if $p : X \to T^2$ is such a bundle then $p^*([T^2]) = 0$. It follows that $L_2(p, \overline{c}) = 0$ for any $c \in T^2$ (base). On the other hand, T^2 is aspherical, so it follows from the main result of [13] that the (abelianized) obstruction $\overline{o}_2(p, \overline{c}) \neq 0$. Again, for *any*

$f : X \to T^2, L_2(p, \overline{c}, f) = 0$. Since T^2 is a Jiang space, it follows from Theorem 6.3 that $p, \overline{c}, f$ are deformable to be coincidence free.

We can now generalize the examples above as follows.

Example 7.3. Let $M = M^{(k-1)n}$ be an even dimensional closed connected oriented manifold and $p : M \to N$ is a fibration over a closed connected oriented n-manifold N with a typical fiber F, a closed connected oriented manifold. Asssume that M is not symplectic, N is aspherical, and is a Jiang-type space as in Theorem 6.3. For any $c \in N$ and any maps $f_3, ..., f_k : M \to N$, $L_2(p, \overline{c}, f_3, ..., f_k) = 0$ so that $p, \overline{c}, f_3, ..., f_k$ are deformable to be coincidence free but $p, \overline{c}$ cannot be homotopic to coincidence free maps.

References

[1] C. Biasi, A. Libardi and T. Monis, The Lefschetz coincidence class of p-maps, *Forum Math.*, **27**(2015), 1717–1728 (DOI 10.1515/forum-2013-0038 online May 2013).

[2] R. Dobreńko, The obstruction to the deformation of a map out of a subspace, *Dissertationes Math. (Rozprawy Mat.)* **295** (1990), 29 pp.

[3] R. Dobreńko and Z. Kucharsky, On the generalization of the Nielsen number, *Fund. Math.* **134** (1990) 1–14.

[4] E. Fadell, On a coincidence theorem of F.B. Fuller, *Pacific J. Math.* **15** (1965), 825–834.

[5] E. Fadell and S. Husseini, Fixed point theory for non simply connected manifolds, *Topology* **20** (1981), 53–92.

[6] H. Geiges, Symplectic structures on T^2-bundles over T^2, *Duke Math. J.* **67** (1992), 539–555.

[7] D. Gonçalves and J. Jezierski, Lefschetz coincidence formula on non-orientable manifolds, *Fund. Math.* **153** (1997), 1–23.

[8] D. Gonçalves, J. Jezierski and P. Wong, Obstruction theory and coincidences in positive codimension, *Acta Math. Sin.* (Engl. Ser.) **22** (2006), no. 5, 1591–1602.

[9] D. Gonçalves and P. Wong, Cohomology of preimages with local coefficients, *Algebr. Geom. Topol.* **6** (2006), 1471–1489.

[10] D. Gonçalves and P. Wong, Obstruction theory and coincidences of maps between nilmanifolds, *Archiv Math.* **84** (2005), 568-576.

[11] D. Gonçalves and P. Wong, Nilmanifolds are Jiang-type spaces for coincidences, *Forum Math.* **13** (2001), 133–141.

[12] D. Gonçalves and P. Wong, Homogeneous spaces in coincidence theory, in: P. Schweitzer, ed., Proceedings of the Tenth Brasilian Topology Conference, São Carlos, Brasil, July 1996. Matemática Contemporânea **13** (1997), 143–158.

[13] D. Gonçalves and P. Wong, Fibrations over aspherical manifolds, *Homology, Homotopy, and Applications* **8** (2006), 257–261.

[14] T. Monis and S. Spiez, Lefschetz coincidence class for several maps, *J. Fixed Point Theory Appl.*, (DOI 10.1007/s11784-015-0266-8) First online: October 31, 2015.

[15] E. Spanier, Duality in topological manifolds, *Colloque de Topologie Tenu á Bruxelles* (Centre Belge de Recherche Mathematiques) (1966), 91-111.

[16] P. Staecker, Nielsen equalizer theory, *Topology Appl.* **158** (2011), 1615–1625; Addendum, ibid, 2256.

[17] W. Thurston, Some simple examples of symplectic manifolds, *Proc. Amer. Math. Soc.* **55**(1976), 467-468.

[18] D. Vendruscolo and P. Wong, Jiang-type theorems for coincidences of maps into homogeneous spaces, *Topol. Methods Nonlinear Anal.*, **31** (2008), 151–160.

[19] G. Whitehead, "Elements of Homotopy Theory," Springer Verlag, New York, 1978.

Dept. de Matemática - UNESP Rio Claro, Av. 24A, 1515 - Bela Vista, Rio Claro-SP, Brasil

E-mail address: tfmonis@rc.unesp.br

Department of Mathematics, Bates College, Lewiston, ME 04240, U.S.A.

E-mail address: pwong@bates.edu